



ENVIRONMENTAL
HEALTH
PERSPECTIVES

<http://www.ehponline.org>

Modeled PFOA Exposure and Coronary Artery Disease, Hypertension, and High Cholesterol in Community and Worker Cohorts

Andrea Winquist and Kyle Steenland

<http://dx.doi.org/10.1289/ehp.1307943>

Received: 30 November 2013

Accepted: 24 September 2014

Advance Publication: 26 September 2014

Modeled PFOA Exposure and Coronary Artery Disease, Hypertension, and High Cholesterol in Community and Worker Cohorts

Andrea Winquist and Kyle Steenland

Environmental Health Department, Rollins School of Public Health, Emory University, Atlanta,
Georgia, USA

Address correspondence to Andrea Winquist, 1518 Clifton Road, NE, Atlanta, GA 30322
USA. Telephone: 404-727-3601. Fax: 404-727-8744. E-mail: awinqui@emory.edu

Running title: PFOA and cardiovascular disease

Acknowledgements: This research is funded by the C8 Class Action Settlement Agreement (Circuit Court of Wood County, West Virginia, USA) between DuPont and Plaintiffs, which resulted from releases into drinking water of the chemical perfluorooctanoic acid (PFOA or C8). Funds are administered by an agency which reports to the court. Our work and conclusions are independent of either party to the lawsuit.

Competing financial interests: The authors declare that they have no competing financial interests.

Abstract

Background: Several previous studies, mostly cross-sectional, have found associations between perfluorooctanoic acid (PFOA) and high cholesterol levels, but studies of hypertension and heart disease have had inconsistent findings.

Objectives: This study examined the association between modeled PFOA exposure and incident hypertension, hypercholesterolemia, and coronary artery disease among workers at a Mid-Ohio Valley chemical plant that used PFOA, and residents of the surrounding community.

Methods: Community- and worker-cohort participants completed surveys during 2008-2011 covering demographics, health-related behaviors, and medical history. Cox proportional hazard models, stratified by birth year, modeled the hazard of each outcome (starting at age 20) as a function of retrospective serum PFOA concentration estimates (generated through fate, transport and exposure modeling), controlling for gender, race, education, smoking, alcohol use, body mass index, and diabetes.

Results: Among 32,254 participants (28,541 community; 3,713 worker), 12,325 reported hypertension with medication, 9,909 reported hypercholesterolemia with medication, and 3,147 reported coronary artery disease (2,550 validated). Hypercholesterolemia incidence increased with increasing cumulative PFOA exposure (sum of yearly serum concentration estimates), most notably among males aged 40-60 years. Compared with the lowest exposure quintile (<142 ng/ml-years), hazard ratios for subsequent quintiles (ng/ml-years: 142-<234; 234-<630; 630-<3579; \geq 3579) were 1.24, 1.17, 1.19 and 1.19 overall and 1.38, 1.32, 1.31, and 1.44 among men aged 40-60 years. There was no apparent association between PFOA exposure and hypertension or coronary artery disease incidence.

Conclusions: Higher PFOA exposure was associated with incident hypercholesterolemia with medication, but not with hypertension or coronary artery disease.

Background

Perfluorooctanoic acid (PFOA) is a man-made 8-carbon perfluorinated compound used as an emulsifier in the manufacture of polytetrafluoroethylene (e.g., Teflon®) and other fluoropolymers (Lau et al. 2007). It is not easily broken down and persists in the environment (Lau et al. 2007). In humans, PFOA has a half-life of approximately 2.3-3.4 years (Bartell et al. 2010; Olsen et al. 2007). In the U.S. National Health and Nutrition Examination Survey (NHANES), PFOA was found in the blood of >99% of US adults during 1999-2008, despite reductions in PFOA emissions since 1999 (Kato et al. 2011). Human exposure in the general population is thought to occur through drinking water, house dust, food (including migration from packaging), and air (Fromme et al. 2009; Lau et al. 2007). Exposures at levels higher than the general population occur in workers and populations living near facilities using PFOA (Lau et al. 2007; Steenland et al. 2010a).

The relationship between PFOA and coronary artery disease is of concern because PFOA has been reported to be associated with several coronary artery disease risk factors, including hypertension (Min et al. 2012), higher serum cholesterol concentrations (Costa et al. 2009; Eriksen et al. 2013; Fisher et al. 2013; Fitz-Simon et al. 2013; Frisbee et al. 2010; Nelson et al. 2010; Olsen et al. 2003; Sakr et al. 2007a; Sakr et al. 2007b; Steenland et al. 2009), higher serum uric acid levels (Geiger et al. 2013; Shankar et al. 2011; Steenland et al. 2010b), and higher serum homocysteine levels (Min et al. 2012). However, studies directly examining the relationship between PFOA and coronary artery disease have had mixed findings. Mortality studies conducted among occupational cohorts (including part of the cohort in this study) have not found associations between PFOA exposure and ischemic heart disease mortality (Leonard et al. 2008; Lundin et al. 2009; Sakr et al. 2009; Steenland and Woskie 2012). One cross-sectional

study of PFOA exposure in the general US population, using NHANES data (1999-2000, 2003-2004 and 2005-2006), also did not find an association with ischemic heart disease (defined as coronary heart disease, angina and/or heart attack) among participants aged ≥ 20 years (Melzer et al. 2010), while a second study using NHANES data (1999-2000 and 2003-2004) did find an association with cardiovascular disease (defined as coronary artery disease, heart attack or stroke) among participants aged ≥ 40 years (Shankar et al. 2012).

The current study is the first to longitudinally examine associations between PFOA exposure and hypercholesterolemia, hypertension, and coronary artery disease incidence among a population with high PFOA exposure. The population included workers at a chemical plant that used PFOA and residents of the surrounding community who were exposed to PFOA through contaminated drinking water. PFOA had been released from the chemical plant into the air, soil and water starting in 1951 (Paustenbach et al. 2007). A class action law suit, filed in 2001-2002, alleged health damage due to PFOA exposure in the surrounding community. A settlement of the lawsuit led to a survey (the “C8 Health Project”), conducted in 2005-2006, of people who had been exposed for at least 12 months (through their primary residence, workplace or school) to water in 6 water districts with PFOA contamination (Frisbee et al. 2009). The C8 Health project survey included questions about medical history (with medical records validation of some conditions), and measurement of PFOA serum concentrations. The lawsuit settlement also formed the C8 Science Panel, a group of epidemiologists charged with determining whether PFOA (also called C8) was linked with any human diseases (c8sciencepanel.org). This study was conducted by the C8 Science Panel.

Methods

Cohort recruitment and survey administration

This study included a community cohort and a worker cohort, which were combined for some analyses and also considered separately. The methods for cohort recruitment and survey administration have been previously described (Winquist et al. 2013). Community cohort participants were recruited among 40,145 C8 Health Project participants who were aged ≥ 20 years and consented to be contacted by the C8 Science Panel for further studies. Worker cohort participants were recruited from an occupational cohort, formed for previous mortality studies (Leonard et al. 2008; Sakr et al. 2009), that included 6,026 people who worked at the chemical plant during 1948-2002 (of whom 2,090 were also in the community cohort). Study participants were asked to complete questionnaires during two survey rounds (2008-2010 and 2010-2011, see Supplemental Material, Figure S1), covering demographics, health-related behaviors, and lifetime personal history of various medical diagnoses. Some participants completed a questionnaire during both survey rounds, and some during only one round. Proxy respondents were sought for people who were deceased or too ill to respond themselves. Participants gave informed consent prior to study participation. This study was approved by the Emory University Institutional Review Board.

At least one study questionnaire was completed during 2008-2011 for 32,712 (81.5%) of the 40,145 people in the community cohort target population (of whom 4,152 were excluded from the community cohort due to working at the plant), and for 4,391 (72.9%) of the 6,026 people in the worker cohort target population. People in the community cohort also in the worker cohort were included only in the worker cohort. Retrospective modeled serum estimates were available for 28,541 (99.9%) community cohort participants and 3,713 (84.6%) worker cohort participants.

Case definitions

On the 2008-2011 surveys, participants were asked whether they had ever been told by a doctor or other health professional that they had high blood pressure, high cholesterol, or heart disease. Women were told to only report hypertension outside of pregnancy. Type of heart disease was also asked; choices included angina, arrhythmia, valve disease, heart attack, coronary artery disease, or 'other' (asked to specify in a free text field). Responses in 'other' fields were reviewed and classified. Reports (through selection of a listed option or through the text field) of angina, coronary atherosclerosis, coronary artery disease (including blocked arteries, bypass surgery, or stents), heart attack, or cardiac ischemia were considered to be reports of coronary artery disease for this analysis. For each reported condition, participants were asked the age at diagnosis. Participants reporting hypertension and high cholesterol were asked whether they were currently taking prescription medication for these conditions. Participants who reported hypertension or high cholesterol without medication during the first survey round were asked again about current medication use during the second survey round. If a proxy respondent reported hypertension or high cholesterol, the proxy was asked whether the person took prescription medication for the condition. Participants reporting heart disease were asked to consent for review of their medical records. Medical records were requested from the identified providers and were reviewed by trained medical record abstractors.

For this analysis, we analyzed incident hypercholesterolemia or hypertension cases reporting current prescription medication on the 2008-2011 surveys. We restricted cases of hypercholesterolemia and hypertension to those with prescription medication use as a way of identifying clinically important conditions. The onset age used in the analysis for these conditions was the date of first diagnosis, not the date of first medication use. People who

reported hypertension or hypercholesterolemia without prescription medication use were excluded from the analysis. People diagnosed with persistent hypercholesterolemia or hypertension typically will take medication over their lifetime. However, we recognize that our case definition could have excluded some people who stopped medication before the survey, or who had been diagnosed but had not yet started medication use by the time of the survey. We think it is likely that these exclusions would be few, and there is no a priori reason why they would be associated with PFOA exposure.

Analyses for heart disease were restricted to cases for whom the medical record documented angina, coronary atherosclerosis, coronary artery disease, heart attack, or cardiac ischemia. We also accepted as valid any self-reported coronary artery disease cases (reported on the 2008-2011 surveys) which had been previously validated in the C8 Health Project survey (2005/2006).

Exposure estimation

Details of the exposure modeling are described elsewhere (Winqvist et al. 2013). Briefly, an environmental fate and transport model was used to generate yearly estimates of PFOA concentrations in local air, surface water, and groundwater (Shin et al. 2011a). These concentrations were used, in combination with survey information relating to residential history, drinking-water sources and water consumption rates, in a residential exposure model to estimate yearly PFOA intake rates (Shin et al. 2011b). Finally, the yearly intake estimates were used in a pharmacokinetic model to generate yearly PFOA serum concentration estimates (Shin et al. 2011b). For people in the worker cohort, job and department-specific yearly PFOA serum concentration estimates were generated in an occupational model based on historical serum PFOA measurements, participants' work histories, and knowledge of plant processes (Woskie et al. 2012). For people in the worker cohort, occupational exposure model estimates were used for

the years when they worked at the plant if they were higher than the residential model estimates; if they were lower, the residential estimates were used. For years after a person stopped working at the plant, serum estimates were decayed 18% per year (based on a half-life of 3.5 years) (Olsen et al. 2007), until they reached a level predicted by the residential model. The Spearman's rank correlation between modeled serum concentration estimates and serum concentrations measured in 2005-2006 (among 30,303 people with serum concentration measurements) was 0.71 (Winquist et al. 2013). For prospective analyses (starting one year after the age at the time of the C8 Health project), estimates were calibrated to the measured serum PFOA concentrations using Bayesian calibration, with measurements weighted more heavily for estimates closer in time to the measurements. Retrospective analyses used un-calibrated estimates due to uncertainty about whether calibration would improve the accuracy of estimates for years far removed from the measurements.

Data analysis

Associations between PFOA serum concentration and the outcomes of interest were examined using Cox proportional hazard models with age as the time scale and time-varying PFOA exposure measures. Retrospective analyses started at the later of age 20 (to consider only adult disease) or the age in 1952 (the year after PFOA production started at the plant). Prospective analyses started at the age one year after the age at C8 Health Project participation (2005/2006) or the age one year after 8/1/2006. Analyses ended at the earliest of the age at diagnosis of the condition of interest, death or the last study questionnaire. Both types of analyses excluded those with a diagnosis of the condition of interest before the analysis start age and those missing a diagnosis age. We excluded people born before 1920 (n=173) because of uncertain reliability of disease reporting in this group. Models for hypercholesterolemia and hypertension excluded

people who reported the condition without current medication (n=5,916 for hypercholesterolemia and 2,470 for hypertension). Models for coronary artery disease excluded people who reported heart disease that was not validated coronary artery disease (597 reported coronary artery disease that was not validated, and 3,728 reported heart disease other than coronary artery disease). Models for coronary artery disease also excluded people without validated coronary artery disease but with coronary artery disease identified in mortality data (n=75). All analyses were done using SAS version 9.2 (SAS Institute, Cary, NC).

Because hypercholesterolemia, hypertension and coronary artery disease are chronic conditions that likely develop over time, our primary exposure metric was a measure of cumulative PFOA exposure, calculated as the sum of all yearly serum concentration estimates for a person through a given age. Because current PFOA serum concentrations could affect these outcomes more acutely, we also used the yearly serum PFOA concentration estimates (at time of case diagnosis or the corresponding age for non-cases) in secondary analyses. Primary analyses considered the exposure by quintiles of the exposure metric, defined among the exposure estimates for cases at their diagnosis age. As a test for trend, we considered the log of the cumulative or yearly serum concentration estimates as a continuous variable.

There was some evidence of violation of the proportional hazards assumption for the cumulative and yearly exposure metrics in retrospective hypercholesterolemia analyses, and in prospective hypertension analyses. Therefore, all analyses were performed both across all ages and separately by age strata (ages 20-39, 40-59, and 60-79 years). All models were stratified by single-year birth year to tightly control for birth year. Models either included a term for the interaction between gender and age or considered genders separately. Models also controlled for years of schooling (not time-varying; <12 years, high school diploma/GED, some college, or

bachelor's degree or higher), race (white vs. non-white or missing; Hispanic ethnicity was not considered because there were only 42 self-reported Hispanics in the study population), smoking (time-varying; current, former, none), smoking duration (time varying), smoking pack years (time-varying linear term created by multiplying the self-reported number of packs smoked per day by the smoking duration to that point), regular alcohol consumption (time-varying; current, former, none), BMI (at time of first study survey; underweight, normal, overweight, obese), and self-reported type 2 diabetes (time-varying according to reported age at diagnosis). Sensitivity analyses that did not control for self-reported type 2 diabetes gave very similar results (data not shown). Family history of coronary artery disease did not change estimates of exposure effect (data not shown) and was not included in models. Models for coronary artery disease did not control for hypercholesterolemia or hypertension because these outcomes were considered to potentially be in the causal pathway.

The primary analyses considered the combined cohorts (88% community, 12% workers). To assess the impact of combining the two cohorts (which likely had different exposure levels, different overall health status, and possibly different exposures to other environmental factors), we considered sensitivity analyses confined to the community cohort. To examine the extent to which results may have been affected by low exposures occurring prior to a person living in the study area, and possible differential migration into the area by disease status, additional sensitivity analyses started at the first age at which each person was known to have qualified for one of the cohorts (by living in the study area for at least one year or working at the plant, referred to as the “qualifying year”). Additional sensitivity analyses considered myocardial infarction as an outcome (a subset of coronary artery disease cases), or were restricted to non-smokers (to consider a group without another strong cardiovascular disease risk factor). To

assess possible effect modification by calendar time, we considered models that ended the cohort follow-up (and hence the analysis) in varying calendar years, in three-year intervals back to 1987 (before peak exposure).

Results

Cohort characteristics

Characteristics of study participants overall and by cohort are shown in Table 1. In the combined cohort, 54% were female, 97% were of white race, 71% were overweight or obese, 52% reported ever smoking, and 14% reported ever having a Type II diabetes diagnosis. The median follow up duration was 32.6 years starting at the age in 1952 or age 20, and 4.4 years starting at the time of the C8 Health Project. The median PFOA serum concentration measured in 2005-2006 was 26.1 ng/ml. Median estimated serum PFOA concentrations (see Supplemental Material, Figure S2) were highest in the late 1990's and early 2000's (Winqvist et al. 2013). Table 2 reports the number of people reporting the outcomes of interest, the number with reported current medication or medical records validation, and the number included in the analyses. The mean age at first diagnosis was 47 years for hypertension among those who reported prescription medication use, 50 years for hypercholesterolemia among those who reported prescription medication use, and 56 years for coronary artery disease among those for whom the condition was validated.

Survival analysis results

Hypertension

In the primary retrospective analysis, combining genders and ages, there was no clear evidence of an association between cumulative PFOA exposure and hypertension (hazard ratios (HRs) and

95% confidence intervals (CI) by increasing quintile: 1.00, 1.10 (1.02, 1.19), 1.10 (1.02, 1.18), 1.05 (0.97, 1.12), 0.98 (0.91, 1.06)) (Figure 1). Results of analysis using the yearly exposure metric were similar (data not shown). In age- and gender-stratified analyses (Figure 1), there was some evidence of increasing hypertension hazard with increasing cumulative or yearly exposure among females in the 20-39 year age group and males in the 40-59 year age group, but trends in hazard across exposure quintiles were not monotonic and tests for trend using the logged exposure metrics were not statistically significant (at $\alpha=0.05$). Prospective analyses starting at the age in 2005-2006, and conditioning on absence of hypertension to that point, also showed no clear evidence of a positive association between cumulative (see Supplemental Material, Figure S3) or yearly PFOA exposure and hypertension, and showed HRs for some quintiles relative to quintile 1 that were below 1.

Retrospective analyses restricted to community cohort members had results similar to the primary analysis (Supplemental Material, Figure S4); trends across PFOA exposure quintiles were not statistically significant (at $\alpha=0.05$). Results of retrospective analysis restricted to non-smokers and analysis starting at the “qualifying age” were similar to the primary analysis (not shown). In retrospective analyses varying the year in which the analysis was ended, an apparent trend (although not completely monotonic) of increasing hypertension hazard with increasing cumulative PFOA exposure quintile was seen for genders and ages combined in analyses ending in 1987; this trend became progressively less clear with longer follow-up after the time of peak exposure (Supplemental Material, Figure S5).

Hypercholesterolemia

The primary retrospective analysis, combining ages and genders, showed an elevated hazard of hypercholesterolemia for exposure quintiles 2-5 relative to quintile 1 (Figure 2). For cumulative

exposure, hazard ratios increased in quintile 2 and remained elevated with little further increase across subsequent quintiles (HRs and 95% CIs by increasing quintile: 1.00, 1.24 (1.15, 1.33), 1.17 (1.09, 1.26), 1.19 (1.11, 1.27), 1.19 (1.11, 1.28); test for trend using log cumulative exposure $p=0.005$). Analyses using the yearly exposure metric also showed an increasing hazard of hypercholesterolemia with increasing quintiles (HRs and 95% CIs by increasing quintile: 1.00, 1.07 (1.01, 1.15), 1.11 (1.04, 1.19), 1.05 (0.99, 1.13), 1.20 (1.12, 1.28); test for trend using log yearly exposure $p<0.001$). In age and gender-specific analyses, the increased hazard for hypercholesterolemia was most pronounced among men aged 40-59 years (HRs and 95% CIs by increasing quintile: cumulative: 1.00, 1.38 (1.21, 1.56), 1.32 (1.17, 1.50), 1.31 (1.16, 1.48), 1.44 (1.28, 1.62), test for trend using log cumulative exposure $p<0.001$; yearly: 1.00, 1.16 (1.03, 1.30), 1.19 (1.05, 1.34), 1.16 (1.03, 1.31), 1.38 (1.23, 1.54), test for trend using log yearly exposure $p<0.001$). In analyses combining ages, tests for interaction between log exposure and gender were statistically significant (cumulative $p=0.048$, yearly $p=0.004$). Tests for interaction with age were also statistically significant (cumulative exposure, genders combined, $p=0.015$, males $p=<0.001$; yearly exposure, males $p=0.032$). Because of these interactions, analyses stratified by gender and age are preferred to the overall analyses. Prospective analyses starting at the age in 2005-2006, and conditioning on absence of hypercholesterolemia to that point, showed no evidence of a positive association between hypercholesterolemia and higher PFOA exposure levels for the cumulative (see Supplemental Material, Figure S3) or yearly exposure metrics with Bayesian calibration (some observed HRs relative to quintile 1 were less than 1), in analyses combining ages and genders, or in analyses by age and gender (data not shown).

In retrospective analyses restricted to community cohort members, there was again evidence of a higher hazard of hypercholesterolemia with higher PFOA exposure in analyses combining

genders and ages (see Supplemental Material, Figure S6). Retrospective analyses restricted to non-smokers (not shown) showed overall patterns of association similar to the primary analysis. Retrospective analyses starting at the “qualifying age” (not shown) also showed similar patterns of association, with significant tests for trend with log cumulative and yearly exposures in analyses combining ages and genders, and for males aged 40-59 years. If analyses were ended in 1987, a strong trend of progressively increasing hazard of hypercholesterolemia with increasing PFOA exposure was seen in analyses combining ages and genders, for both the cumulative (log-linear trend test $p < 0.001$) and yearly metrics (log-linear trend test $p < 0.001$) (see Supplemental Material, Figure S7 for yearly metric results). With increasing analysis end years, the pattern of increasing hypercholesterolemia hazard with increasing PFOA exposure became progressively weaker for both exposure metrics.

Coronary artery disease

In the primary retrospective analysis there was no clear association between PFOA exposure and coronary artery disease (Figure 3) (combining genders and ages, HRs and 95% CIs by increasing quintile: cumulative 1.00, 1.26 (1.10, 1.45), 1.17 (1.02, 1.35), 0.99 (0.86, 1.14), 1.07 (0.93, 1.23); yearly 1.00, 1.02 (0.90, 1.16), 1.06 (0.93, 1.21), 0.95 (0.84, 1.09), 0.97 (0.85, 1.11)). Among males aged 20-39 years, there appeared to be higher hazards of coronary artery disease in quintiles 2-5 relative to quintile 1, but confidence intervals were wide for this age group, and there was no significant trend (log linear test for trend, cumulative exposure $p = 0.64$). Prospective analyses also showed no evidence of increasing hazards for coronary artery disease with increasing PFOA exposure, and showed HRs for quintiles 2-5 relative to quintile 1 that were often below 1 (see Supplemental Material, Figure S3).

In retrospective analyses restricted to community cohort members, patterns of associations were similar to those seen for the combined cohorts, with no clear associations (Supplemental Material, Figure S8). Analyses restricted to non-smokers and analyses starting at the “qualifying age” (not shown) were also similar. Analyses varying the year in which the analysis ended did not show clear monotonic trends across exposure quintiles, but had a suggestion of an association with coronary artery disease for cumulative and yearly PFOA exposure (for the genders combined and among men) when the analysis was ended in years closer to peak PFOA exposures (see Supplemental Material, Figure S9 for yearly exposure results among men) (for analyses ending in 1987, log linear tests for trend: ages and genders combined, cumulative $p=0.07$, yearly $p=0.03$; men, ages combined cumulative $p=0.11$, yearly $p=0.061$). Primary retrospective analysis for myocardial infarction also showed patterns of associations that were similar to those seen for coronary artery disease overall (data not shown).

Discussion

In summary, there was little evidence of an increasing hazard of hypertension with increasing PFOA exposure, but there was evidence of an increasing hazard of hypercholesterolemia with increasing PFOA exposure, (most evident among males aged 40-59 years). Despite the observed association with hypercholesterolemia, there was no clear association between PFOA exposure and coronary artery disease. Although there was a suggestion of an association with coronary artery disease among males aged 20-39, confidence intervals were wide for that group.

Our finding of an association between PFOA exposure and hypercholesterolemia is consistent with the findings of several previous studies. In humans, statistically significant positive associations between PFOA and serum cholesterol levels have been found in 10 studies

(primarily cross sectional) of people occupationally and non-occupationally exposed to PFOA (see Introduction); positive associations that were not statistically significant were found in four additional studies (Emmett et al. 2006; Olsen et al. 2000; Olsen and Zobel 2007; Wang et al. 2012). A longitudinal study of changes in serum cholesterol levels in relation to changes in serum PFOA levels, conducted in a subset of the same population considered in this study, found a small but significant association between decreases in serum PFOA concentrations and decreases in serum cholesterol levels (Fitz-Simon et al. 2013). A second longitudinal study of 179 workers involved with demolition of two former PFOA and PFOS manufacturing facilities (in Minnesota and Alabama), with a mean follow up of 164 days, did not find an association between changes in PFOA serum concentrations and changes in total cholesterol levels between a baseline assessment (before work on demolition) and an end-of-project assessment, either among the group overall or among the subgroup of 120 workers with baseline PFOA levels <15 ng/mL and PFOS levels <50 ng.ml (Olsen et al. 2012).

The evidence for associations between PFOA and either hypertension or coronary artery disease is sparse. One cross-sectional study of general population PFOA exposures, using data from NHANES, found an association between PFOA and systolic blood pressure and hypertension (Min et al. 2012). There has been little evidence of an association between PFOA exposure and coronary artery disease in previous occupational mortality studies of two cohorts (one of which considered the worker cohort in this study) (Lundin et al. 2009; Steenland and Woskie 2012). One cross-sectional study of PFOA exposure in the general US population, using data from NHANES (1999-2000, 2003-2004, and 2005-2006) among people aged ≥ 20 years, also did not find an association between PFOA exposure and ischemic heart disease (defined as coronary artery disease, angina, and/or heart attack) (Melzer et al. 2010), but a second cross-sectional

study of PFOA exposure using NHANES data (1999-2000 and 2003-2004), among people aged ≥ 40 years did find a statistically significant association with cardiovascular disease (defined as coronary artery disease, heart attack or stroke) (Shankar et al. 2012).

The lack of an observed association between PFOA exposure and coronary artery disease in our study population may seem puzzling in light of the evidence for an association with hypercholesterolemia. As expected, time-varying hypertension and hypercholesterolemia were both strongly associated with an increased hazard of coronary artery disease in our study, after controlling for the other variables in our models (results not shown). Nevertheless, effects of strong risk factors for coronary artery disease, such as smoking, could mask an indirect effect of PFOA operating through hypercholesterolemia as an intermediate outcome. However, PFOA also was not associated with coronary artery disease in a sensitivity analysis restricted to non-smokers. In addition, treatment of hypercholesterolemia among our study participants might have mitigated potential effects of PFOA-associated hypercholesterolemia on coronary artery disease.

Strengths of this study include examination of a large cohort with exposure estimates going back to birth or 1951, consideration of disease incidence (rather than mortality or prevalence), and validation of coronary artery disease diagnoses. This study also has several limitations. To be in the community cohort, a person had to be alive in 2005-2006, so some people who developed coronary artery disease may have been unavailable for enrollment in our study. Although workers did not have to be alive in 2005-2006, proxies were difficult to find for deceased workers. The impact on the analysis of loss of people who died of coronary artery disease is uncertain, since it is unclear whether this loss would be related to PFOA exposure. This study is also limited by the fact that exposure had been ongoing in this population for many years. If

susceptibility to effects of PFOA on our outcomes varied in the population, this could result in progressively decreasing HRs with increasing follow-up time due to progressively decreasing susceptibility of the remaining population (Applebaum et al. 2011; Hernan 2010). We saw some evidence that this may have affected our results in analyses that ended in years closer to peak exposures. This issue may have particularly affected our prospective analyses (which were conditional on lack of disease prior to 2005-2006), and could explain prospective HRs relative to quintile 1 that were below 1.

Our primary analyses combined the community and worker cohorts to examine associations between PFOA and our outcomes across a wide exposure range. However, because workers had the highest exposures and may have been different from non-workers in important ways, we considered sensitivity analyses restricted to non-workers. Results of those analyses were generally similar to analyses including workers, suggesting that inclusion of workers did not lead to substantial bias in our results.

Finally, there is a potential for misclassification of disease, exposure, and covariates due to inaccuracies in modeled exposure estimates and in self-reporting of disease and covariates. Any exposure misclassification would be expected to be independent of disease status and would likely, but not certainly, bias results toward the null. We sought to minimize misclassification of disease status by requiring prescription medication use for hypercholesterolemia and hypertension, and medical records validation for coronary artery disease (to increase disease classification specificity); and by excluding from the analysis people who had some indication of disease, but did not meet the case definitions.

In conclusion, our study provided additional evidence supporting an association between PFOA exposure and hypercholesterolemia. However, we did not find strong evidence of an association between PFOA exposure and hypertension or coronary artery disease.

References

- Applebaum KM, Malloy EJ, Eisen EA. 2011. Left Truncation, Susceptibility, and Bias in Occupational Cohort Studies. *Epidemiology* 22:599-606.
- Bartell SM, Calafat AM, Lyu C, Kato K, Ryan PB, Steenland K. 2010. Rate of Decline in Serum PFOA Concentrations after Granular Activated Carbon Filtration at Two Public Water Systems in Ohio and West Virginia. *Environ Health Persp* 118:222-228.
- Costa G, Sartori S, Consonni D. 2009. Thirty Years of Medical Surveillance in Perfluorooctanoic Acid Production Workers. *J Occup Environ Med* 51:364-372.
- Emmett EA, Zhang H, Shofer FS, Freeman D, Rodway NV, Desai C, et al. 2006. Community exposure to perfluorooctanoate: relationships between serum levels and certain health parameters. *J Occup Environ Med* 48:771-779.
- Eriksen KT, Raaschou-Nielsen O, McLaughlin JK, Lipworth L, Tjønneland A, Overvad K, et al. 2013. Association between Plasma PFOA and PFOS Levels and Total Cholesterol in a Middle-Aged Danish Population. *Plos One* 8(2).
- Fisher M, Arbuckle TE, Wade M, Haines DA. 2013. Do perfluoroalkyl substances affect metabolic function and plasma lipids?-Analysis of the 2007-2009, Canadian Health Measures Survey (CHMS) Cycle 1. *Environ Res* 121:95-103.
- Fitz-Simon N, Fletcher T, Luster MI, Steenland K, Calafat AM, Kato K, et al. 2013. Reductions in Serum Lipids with a 4-year Decline in Serum Perfluorooctanoic Acid and Perfluorooctanesulfonic Acid. *Epidemiology* 24:569-576.
- Frisbee SJ, Brooks AP, Jr., Maher A, Flensburg P, Arnold S, Fletcher T, et al. 2009. The C8 health project: design, methods, and participants. *Environ Health Perspect* 117:1873-1882.
- Frisbee SJ, Shankar A, Knox SS, Steenland K, Savitz DA, Fletcher T, et al. 2010. Perfluorooctanoic acid, perfluorooctanesulfonate, and serum lipids in children and adolescents: results from the C8 Health Project. *Arch Pediatr Adolesc Med* 164:860-869.
- Fromme H, Tittlemier SA, Volkel W, Wilhelm M, Twardella D. 2009. Perfluorinated compounds - Exposure assessment for the general population in western countries. *Int J Hyg Envir Health* 212:239-270.
- Geiger SD, Xiao J, Shankar A. 2013. Positive Association Between Perfluoroalkyl Chemicals and Hyperuricemia in Children. *Am J Epidemiology* 177:1255-1262.
- Hernan MA. 2010. The Hazards of Hazard Ratios. *Epidemiology* 21:13-15.

- Kato K, Wong LY, Jia LT, Kuklennyik Z, Calafat AM. 2011. Trends in Exposure to Polyfluoroalkyl Chemicals in the US Population: 1999-2008. *Environ Sci Technol* 45:8037-8045.
- Lau C, Anitole K, Hodes C, Lai D, Pfahles-Hutchens A, Seed J. 2007. Perfluoroalkyl acids: a review of monitoring and toxicological findings. *Toxicol Sci* 99:366-394.
- Leonard RC, Kreckmann KH, Sakr CJ, Symons JM. 2008. Retrospective cohort mortality study of workers in a polymer production plant including a reference population of regional workers. *Ann Epidemiol* 18:15-22.
- Lundin JI, Alexander BH, Olsen GW, Church TR. 2009. Ammonium Perfluorooctanoate Production and Occupational Mortality. *Epidemiology* 20:921-928.
- Melzer D, Rice N, Depledge MH, Henley WE, Galloway TS. 2010. Association between Serum Perfluorooctanoic Acid (PFOA) and Thyroid Disease in the US National Health and Nutrition Examination Survey. *Environ Health Persp* 118:686-692.
- Min JY, Lee KJ, Park JB, Min KB. 2012. Perfluorooctanoic acid exposure is associated with elevated homocysteine and hypertension in US adults. *Occup Environ Med* 69:658-662.
- Nelson JW, Hatch EE, Webster TF. 2010. Exposure to Polyfluoroalkyl Chemicals and Cholesterol, Body Weight, and Insulin Resistance in the General US Population. *Environ Health Persp* 118:197-202.
- Olsen GW, Burris JM, Burlew MM, Mandel JH. 2003. Epidemiologic assessment of worker serum perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) concentrations and medical surveillance examinations. *J Occup Environ Med* 45:260-270.
- Olsen GW, Burris JM, Burlew MM, Mandel JH. 2000. Plasma cholecystokinin and hepatic enzymes, cholesterol and lipoproteins in ammonium perfluorooctanoate production workers. *Drug Chem Toxicol* 23:603-620.
- Olsen GW, Burris JM, Ehresman DJ, Froehlich JW, Seacat AM, Butenhoff JL, et al. 2007. Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired fluorochemical production workers. *Environ Health Persp* 115:1298-1305.
- Olsen GW, Ehresman DJ, Buehrer BD, Gibson BA, Butenhoff JL, Zobel LR. 2012. Longitudinal Assessment of Lipid and Hepatic Clinical Parameters in Workers Involved With the Demolition of Perfluoroalkyl Manufacturing Facilities. *J Occup Environ Med* 54:974-983.

- Olsen GW, Zobel LR. 2007. Assessment of lipid, hepatic, and thyroid parameters with serum perfluorooctanoate (PFOA) concentrations in fluorochemical production workers. *Int Arch Occup Environ Health* 81:231-246.
- Paustenbach DJ, Panko JM, Scott PK, Unice KM. 2007. A methodology for estimating human exposure to perfluorooctanoic acid (PFOA): a retrospective exposure assessment of a community (1951-2003). *J Toxicol Environ Health A* 70:28-57.
- Sakr CJ, Kreckmann KH, Green JW, Gillies PJ, Reynolds JL, Leonard RC. 2007a. Cross-sectional study of lipids and liver enzymes related to a serum biomarker of exposure (ammonia perfluorooctanoate or APFO) as part of a general health survey in a cohort of occupational exposed workers. *J Occup Environ Med* 49:1086-1096.
- Sakr CJ, Leonard RC, Kreckmann KH, Slade MD, Cullen MR. 2007b. Longitudinal study of serum lipids and liver enzymes in workers with occupational exposure to ammonium perfluorooctanoate. *J Occup Environ Med* 49:872-879.
- Sakr CJ, Symons JM, Kreckmann KH, Leonard RC. 2009. Ischaemic heart disease mortality study among workers with occupational exposure to ammonium perfluorooctanoate. *Occup Environ Med* 66:699-703.
- Shankar A, Xiao J, Ducatman A. 2011. Perfluoroalkyl chemicals and elevated serum uric acid in US adults. *Clin Epidemiol* 3:251-258.
- Shankar A, Xiao J, Ducatman A. 2012. Perfluorooctanoic Acid and Cardiovascular Disease in US Adults. *Arch Intern Med* 172:1397-1403.
- Shin HM, Vieira VM, Ryan PB, Detwiler R, Sanders B, Steenland K, et al. 2011a. Environmental Fate and Transport Modeling for Perfluorooctanoic Acid Emitted from the Washington Works Facility in West Virginia. *Environ Sci Technol* 45:1435-1442.
- Shin HM, Vieira VM, Ryan PB, Steenland K, Bartell SM. 2011b. Retrospective Exposure Estimation and Predicted versus Observed Serum Perfluorooctanoic Acid Concentrations for Participants in the C8 Health Project. *Environ Health Persp* 119:1760-1765.
- Steenland K, Fletcher T, Savitz DA. 2010a. Epidemiologic evidence on the health effects of perfluorooctanoic acid (PFOA). *Environ Health Perspect* 118:1100-1108.
- Steenland K, Tinker S, Frisbee S, Ducatman A, Vaccarino V. 2009. Association of perfluorooctanoic acid and perfluorooctane sulfonate with serum lipids among adults living near a chemical plant. *Am J Epidemiol* 170:1268-1278.

- Steenland K, Tinker S, Shankar A, Ducatman A. 2010b. Association of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) with uric acid among adults with elevated community exposure to PFOA. *Environ Health Perspect* 118:229-233.
- Steenland K, Woskie S. 2012. Cohort Mortality Study of Workers Exposed to Perfluorooctanoic Acid. *Am J Epidemiol* 176:909-917.
- Wang JS, Zhang YT, Zhang W, Jin YH, Dai JY. 2012. Association of Perfluorooctanoic Acid with HDL Cholesterol and Circulating miR-26b and miR-199-3p in Workers of a Fluorochemical Plant and Nearby Residents. *Environ Sci Technol* 46:9274-9281.
- Winqvist A, Lally C, Shin HM, Steenland K. 2013. Design, methods, and population for a study of PFOA health effects among highly exposed mid-Ohio valley community residents and workers. *Environ Health Perspect* 121:893-899.
- Woskie SR, Gore R, Steenland K. 2012. Retrospective Exposure Assessment of Perfluorooctanoic Acid Serum Concentrations at a Fluoropolymer Manufacturing Plant. *Ann Occup Hyg* 56:1025-1037.

Table 1. Population characteristics.

Characteristic	Community Cohort Only (N=28,541)	Worker Cohort Only (N=3,713)	Combined Cohort (N=32,254)
Female	16,602 (58%)	758 (20%)	17,360 (54%)
Birth year			
Before 1920	114 (0.4%)	59 (2%)	173 (0.5%)
1920-1939	3,742 (13%)	727 (20%)	4,469 (14%)
1940-1959	11,547 (40%)	1,814 (49%)	13,361 (41%)
≥1960	13,138 (46%)	1,113 (30%)	14,251 (44%)
Education (4 missing)			
<High school	3,026 (11%)	37 (1%)	3,063 (9%)
High school diploma or GED	11,706 (41%)	1,265 (34%)	12,971 (40%)
Some College	9,441 (33%)	1,081 (29%)	10,522 (33%)
Bachelor's degree or higher	4,366 (15%)	1,328 (36%)	5,694 (18%)
Race (295 missing)			
White	27,901 (98%)	3,284 (88%)	31,185 (97%)
Other	640 (2%)	134 (4%)	774 (2%)
BMI at time of survey (79 missing)			
Underweight (BMI <18.5 kg/m ³)	414 (1%)	38 (0.1%)	452 (1%)
Normal weight (BMI 18.5-<25.0 kg/m ³)	7,693 (27%)	972 (26%)	8,665 (27%)
Overweight (BMI 25.0-<30.0 kg/m ³)	9,689 (34%)	1,618 (44%)	11,307 (35%)
Obese (BMI ≥30.0 kg/m ³)	10,694 (37%)	1,057 (28%)	11,751 (36%)
Smoking			
Never smoked	13,527 (47%)	1,989 (54%)	15,516 (48%)
Smoked and quit	8,899 (31%)	1,297 (35%)	10,196 (32%)
Smoked and did not quit	6,115 (21%)	427 (12%)	6,542 (20%)
Regular alcohol consumption (74 missing)			
Never	17,011 (60%)	1,683 (45%)	18,694 (58%)
Yes and quit	4,105 (14%)	535 (14%)	4,640 (14%)
Yes and did not quit	7,360 (26%)	1,486 (40%)	8,846 (27%)
Any self-reported type II diabetes	3,912	522	4,434 (13.8%)
Serum PFOA concentration measurement from the C8 Health Project (2005-2006) (ng/ml)			
Mean	70.9	324.6	86.6
Standard deviation	151.2	920.6	278.9
25th percentile	12.3	55.9	12.8
Median	24.2	112.7	26.1
75th percentile	58.9	256.2	68.1
Number with measurements	28,422	1,881	30,303

Table 2. Cardiovascular outcome frequencies.

Outcome	# Reporting outcome	# Reporting current medication or validated^a (% of reported)	Number included in the primary retrospective analysis^b	Number included in the prospective analysis^c
Hypertension	14,795	12,325 (83%)	11,798	2,226
Hypercholesterolemia	15,825	9,909 (63%)	9,653	1,825
Coronary artery disease	3,147	2,550 (81.0%) ^a	2,468	515

^aValidation was done only for coronary artery disease. Current medication includes self-reported use of current medication at the time of either survey round, or any medication use reported by a proxy. ^bHad age at diagnosis and all other model covariates. ^cHad age at diagnosis and all other model covariates, and age at diagnosis was after the age at the time of the C8 Health Project.

Figure Legends

Figure 1. Hazard ratios and 95% CI for hypertension in the primary retrospective analysis for the combined cohorts, cumulative exposure. Quintile cut points (in $\mu\text{g}/\text{ml}\cdot\text{yr}$) were: <0.111 , $0.111-<0.191$, $0.191-<0.471$, $0.471-<2.763$, ≥ 2.763 . The analysis included 11,798 cases of self-reported hypertension with medication. Models were stratified by single-year birth year and were either stratified by gender or controlled for gender and the interaction between gender and age. Models also controlled for years of schooling (not time-varying; <12 years, high school diploma/GED, some college, or bachelor's degree or higher), race (white vs. non-white or missing), smoking (time-varying; current, former, none), smoking duration (time varying), smoking pack years (time-varying linear term created by multiplying the self-reported number of packs smoked per day by the smoking duration to that point), regular alcohol consumption (time-varying; current, former, none), BMI (at time of first study survey; underweight, normal, overweight, obese), and self-reported type 2 diabetes (time-varying according to reported age at diagnosis).

Figure 2. Hazard ratios and 95% CI for hypercholesterolemia in the primary retrospective analysis for the combined cohorts, cumulative exposure. Quintile cut points (in $\mu\text{g}/\text{ml}\cdot\text{yr}$) were: <0.142 , $0.142-<0.234$, $0.234-<0.630$, $0.630-<3.579$, ≥ 3.579 . The analysis included 9,653 cases of self-reported hypercholesterolemia with medication. Models were stratified by single-year birth year and were either stratified by gender or controlled for gender and the interaction between gender and age. Models also controlled for years of schooling (not time-varying; <12 years, high school diploma/GED, some college, or bachelor's degree or higher), race (white vs. non-white or missing), smoking (time-varying; current, former, none), smoking duration (time varying), smoking pack years (time-varying linear term created by multiplying the self-reported number of packs smoked per day by the smoking duration to that point), regular alcohol consumption (time-

varying; current, former, none), BMI (at time of first study survey; underweight, normal, overweight, obese), and self-reported type 2 diabetes (time-varying according to reported age at diagnosis).

Figure 3. Hazard ratios and 95% CI for coronary artery disease in the primary retrospective analysis for the combined cohorts, cumulative exposure. Quintile cut points (in $\mu\text{g/ml}\cdot\text{yr}$) were: <0.147 , $0.147-<0.248$, $0.248-<0.717$, $0.717-<5.058$, ≥ 5.058 . The analysis included 2,468 cases of validated coronary artery disease. Models were stratified by single-year birth year and were either stratified by gender or controlled for gender and the interaction between gender and age. Models also controlled for years of schooling (not time-varying; <12 years, high school diploma/GED, some college, or bachelor's degree or higher), race (white vs. non-white or missing), smoking (time-varying; current, former, none), smoking duration (time varying), smoking pack years (time-varying linear term created by multiplying the self-reported number of packs smoked per day by the smoking duration to that point), regular alcohol consumption (time-varying; current, former, none), BMI (at time of first study survey; underweight, normal, overweight, obese), and self-reported type 2 diabetes (time-varying according to reported age at diagnosis).

Figure 1.




Figure 2.




Figure 3.

